

**DTIC FILE COPY**



2

**NAVAL MEDICAL RESEARCH INSTITUTE**

Bethesda, MD 20889-5055

NMRI 90-110

November 1990

**AD-A230 228**

**CAFFEINE ATTENUATES THE AFTERDROP IN RECTAL TEMPERATURE AFTER  
MILD COOLING**

**DTIC**  
**ELECTE**  
**JAN 04 1991**  
**S D**

**S. T. Ahlers**  
**R. L. Hesslink**  
**S. B. Lewis**  
**J. R. Thomas**

**Naval Medical Research  
and Development Command  
Bethesda, Maryland 20889-5055**

**Department of the Navy  
Naval Medical Command  
Washington, DC 20372-5210**

**Approved for public release;  
distribution is unlimited**

**01-1 2 062**

## **NOTICES**

The opinions and assertions contained herein are the private ones of the writer and are not to be construed as official or reflecting the views of the naval service at large.

When U. S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever, and the fact that the Government may have formulated, furnished or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Please do not request copies of this report from the Naval Medical Research Institute. Additional copies may be purchased from:

**National Technical Information Service  
5285 Port Royal Road  
Springfield, Virginia 22161**

Federal Government agencies and their contractors registered with the Defense Technical Information Center should direct requests for copies of this report to:

**Defense Technical Information Center  
Cameron Station  
Alexandria, Virginia 22304-6145**

## **TECHNICAL REVIEW AND APPROVAL NMRI 90-110**

The experiments reported herein were conducted according to the principles set forth in the current edition of the "Guide for the Care and Use of Laboratory Animals," Institute of Laboratory Animal Resources, National Research Council.

This technical report has been reviewed by the NMRI scientific and public affairs staff and is approved for publication. It is releasable to the National Technical Information Service where it will be available to the general public, including foreign nations.

**LARRY W. LAUGHLIN  
CAPT, MC, USN  
Commanding Officer  
Naval Medical Research Institute**

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCL		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMR1		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research Institute	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
6c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20889-5055		7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, DC 20372-5120	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research & Development Command	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) 8901 Wisconsin Avenue Bethesda, MD 20814-5044		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
		63706N	M00095.004 1008 DN246556
11. TITLE (Include Security Classification) Caffeine attenuates the afterdrop in rectal temperature after mild cooling			
12. PERSONAL AUTHOR(S) Stephen T. Ahlers, Robert L. Hesslink, Stephen B. Lewis, and John R. Thomas			
13a. TYPE OF REPORT Technical Report	13b. TIME COVERED FROM 10/85 TO 9/93	14. DATE OF REPORT (Year, Month, Day) 1990 November 26	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)  Vasoconstriction, Rewarming, Catecholamines	
FIELD	GROUP SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Previous research has shown that subjects who experience immersion hypothermia often show a continued decrease in rectal temperature (afterdrop) during the initial phase of rewarming. Because caffeine, the principal constituent of coffee, influences thermoregulatory and cardiovascular responses and is occasionally administered to individuals upon removal from cold exposure, we examined whether caffeine might influence rewarming after mild cooling. Seven male subjects were given either warm decaffeinated coffee containing 3.5 mg/kg caffeine or warm water after their rectal temperature was reduced by 0.5°C via immersion in cold (13°C) water. Between 20 and 40 minutes (min) after removal from cold water a 0.8°C afterdrop in rectal temperature occurred in the noncaffeine control condition, while the condition			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Regina E. Hunt, Command Editor		22b. TELEPHONE (Include Area Code) (301) 295-0198	22c. OFFICE SYMBOL SD/RSD/NMRI

9. administered caffeine showed a mean afterdrop of  $0.5^{\circ}\text{C}$ . After 120 min, the caffeine condition had rewarmed back to the pre-afterdrop criterion, while the control condition was still 0.5 below the criterion level. These data indicate that caffeine significantly attenuated the magnitude of the post-immersion afterdrop in rectal temperature.

35 ←

# TABLE OF CONTENTS

	<u>Page No.</u>
Acknowledgements	iv
Introduction	1
Methods	2
Results	4
Discussion	9
References	12

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

### ACKNOWLEDGEMENTS

The authors gratefully acknowledge Dr. L. Homer, Mr..R. Robertson, HM2 L. Allen, HM2 V. Harmon, HM1 P. Wilson, and Ms. P. Mullinix for their valuable contributions to this project. This report was supported by the Naval Medical Research and Development Command Work unit #63706N.M00095.004.1008. Opinions and assertions contained herein are the private views of the authors and are not to be construed as official or reflecting the views of the Department of Defense, the Department of Navy, or the Naval Service at large.

## INTRODUCTION

After removal from extended exposure in cold water, which induces core hypothermia, body temperature often continues to decline (afterdrop) during the initial phase of rewarming. If the temperature becomes low enough, the condition can be life threatening (1). Although the mechanisms underlying this phenomenon remain to be precisely elucidated, current hypotheses suggest that post-immersion afterdrop stems from a continuation of both conductive heat loss through tissue (2) and convective heat loss, resulting from the perfusion of cooled blood from the periphery into the deep core (3).

In the field, coffee is often given to individuals exposed to cold as a means of introducing warm fluids. However, the principal constituent of coffee, caffeine, produces a number of cardiovascular and metabolic effects that could influence the physiological response to cold. Oral caffeine, when given alone or in coffee, increases peripheral vasoconstriction, vascular resistance, systemic blood pressure, and heart rate (4, 5, 6, 7); caffeine has also been shown to decrease peripheral skin temperature (7). These physiological effects, which collectively result from increased heart rate and vascular resistance, are thought to be mediated by antagonism of adenosine (8), which increases the systemic release of the adrenergic catecholamines, epinephrine and norepinephrine (6, 9, 10).

Given its pervasive physiological effects as well as its use in the field, we examined whether caffeine might influence the

rate of rewarming after moderate hypothermia was induced by immersion in cold water.

## METHODS

### Subjects:

Seven adult male volunteers in good general health served as subjects. Once they were briefed on the risks and benefits of the study, all participants provided signed informed consent. Medical coverage was provided for all subjects. Individual subject characteristics are provided in Table 1.

All subjects were instructed to abstain from any caffeinated substance for 24 hours prior to the experimental sessions. On two occasions, separated by at least one week, each subject arrived at the laboratory after a 12 hour overnight fast and was

**Table 1. Subject Characteristics**

<u>SUBJECT</u>	<u>% FAT</u>	<u>SURFACE AREA (m<sup>2</sup>)</u>	<u>AGE</u>	<u>Ht (cm)</u>	<u>Wt (Kg)</u>	<u>IM1</u>	<u>IM2</u>
1	10	1.95	23	180.3	75.9	30	35
2	11	1.77	25	172.2	64.5	28	13
3	10	1.83	29	172.7	70.5	22	30
4	11	1.87	26	177.8	70.5	20	30
5	20	2.19	30	186.7	94.0	40	35
6	17	2.06	30	177.8	88.0	50	55
7	14	1.89	27	177.8	71.8	45	36
MEAN	<u>13</u>	<u>1.94</u>	<u>27</u>	<u>177.9</u>	<u>76.6</u>	<u>34</u>	<u>33</u>
SD	3.6	0.13	2.5	4.52	9.8	10.7	10.7

Percent body fat (% fat) was determined by hydrostatic weighing. Surface area was calculated according to the method of DuBois and DuBois (11). Immersion times are expressed in minutes.



weighed. Subjects were instrumented with a YSI rectal probe, inserted 15 cm past the anal sphincter. Additional YSI 700 series thermistors (Yellow Springs, OH) were taped bilaterally to the index fingers, triceps, quadriceps, and great toes. Heart rate was monitored with a standard 3-lead EKG monitor. An 18 gauge indwelling catheter was placed into the antecubital vein of the non-immersed arm and was kept patent with heparinized saline. Temperature and heart rate data were recorded manually at 10-minute intervals throughout the experimental procedure.

Wearing swimming trunks only, subjects remained seated for 30 minutes in order that stable baseline measurements could be obtained. Thereafter, each subject was immersed up to the shoulder level in 13°C circulating water until his rectal temperature decreased by 0.5°C. Upon reaching the temperature criterion, subjects were immediately removed from the immersion bath, towel dried, and allowed to rewarm for 160 minutes while seated upright; ambient air temperature in the room was kept constant between 24-26°C. Within one minute after removal from the immersion bath, each subject was orally administered either caffeine (3.5 mg/kg) mixed in 500 ml of warm (49-50°C) decaffeinated coffee or warm water (control); presentation of these conditions was counterbalanced between sessions for all subjects. The amount of caffeine was approximately the amount contained in two to three cups of brewed coffee.

Blood samples were taken just before cold water immersion, at time zero when subjects reached the criterion temperature, and

at 10 and 20 minutes after removal from the immersion bath. Thereafter, samples were drawn at 20-minute intervals in the rewarming period. Samples were drawn into cooled tubes containing 14.3 USP units/ml of heparin and were immediately cooled. The blood was subsequently separated into plasma by centrifugation at 3000 rpm for 11 minutes at 4°C. Following centrifugation, 1 ml aliquots of plasma were pipetted into tubes containing 7.5 nM EGTA and 5.0 nM glutathione in deionized water at PH 6.8; this preparation was frozen and maintained at -80°C for subsequent analysis. Thawed extracts eluted from alumina were analyzed for epinephrine and norepinephrine by use of high pressure liquid chromatography with electrochemical detection; 3,4-dihydroxybenzylamine was used as the internal standard. Interassay variation with this method of catecholamine analysis was determined to be less than 10%.

## RESULTS

### Cooling:


The time taken to reach a 0.5°C reduction in rectal temperature on each exposure is shown in the right-hand portion of Table 1. There were no significant differences in the length of time to reach this temperature criterion between the two experimental conditions, nor was the rate of cooling different between the first and second exposure ( $p$ 's > .05).

### Rewarming:


Administration of 3.5 mg/kg caffeine within one minute after subjects were removed from the cold water significantly modified

rectal temperature (Fig. 1) during the 160-min rewarming phase ( $F = 2.34$ ,  $p < .008$ ) such that the rectal temperature of the caffeine condition was significantly higher throughout the rewarming period. Analysis of the rate of temperature decline for each subject in the first 40 minutes of the rewarming phase indicated no significant differences between the caffeine (slope = 1.21) and control condition [(slope = 1.20),  $p > .05$ ]. At 40 minutes into the rewarming period, rectal temperature was no longer observed to decline any further; however, the rectal temperature of the caffeine condition was approximately  $0.3^{\circ}\text{C}$  above the mean temperature of the control condition. Statistical analysis of the 50 to 160-min time period when temperatures were increasing also revealed nonsignificant differences between the caffeine (slope = 0.315) and control [(slope = 0.311),  $p > .05$ ] conditions. This finding indicates that there were no systematic differences in the rate of rewarming after subjects had reached an asymptotic temperature level.

Analysis of the weighted skin temperatures indicated that caffeine did not significantly modify rewarming of peripheral skin temperature (Fig. 2). The temperature of the skin fell to approximately  $14^{\circ}\text{C}$  during immersion in the cold water but increased rapidly to near basal levels within the first 40 min of the rewarming period. None of the individual thermistor sites were significantly different between the treatment conditions, thus indicating that caffeine had no discernable effect on peripheral temperature during rewarming after mild hypothermia.




Figs. 1 and 2. Mean rectal and skin temperature, respectively, at baseline (B) and during the 160-min rewarming period.




Similarly, caffeine had no effect on plasma norepinephrine (Fig. 3) during the rewarming phase in five of the seven subjects from whom complete blood samples were obtained. Subjects' norepinephrine levels were, however, substantially elevated both during and immediately after exposure to the cold. Removal from the cold water resulted in a further increase of plasma norepinephrine in the initial 10 min of the rewarming period. Thereafter, norepinephrine levels steadily declined throughout the remainder of the 160-min session.

Plasma epinephrine was substantially elevated during the post-immersion rewarming phase for the caffeine condition; however, this occurrence did not reach statistical significance above the control values ( $F = 2.04$ ,  $p = .07$ ) (Fig. 4), perhaps because only samples from five of the seven subjects were obtained. What is clear, however, is that after cold water immersion, subjects' epinephrine levels decreased substantially during the initial 20 minutes of the rewarming period and remained at a sustained low level during the rest of the session. Caffeine appeared to block the post-immersion reduction in epinephrine levels. A similar pattern occurred with the heart rate measure; however, in this case a significant ( $F = 2.57$ ,  $p < .025$ ) difference in heart rate (Fig. 5) was observed. The pattern of the heart rate appeared to parallel the observed changes in plasma epinephrine in that, without caffeine, heart rate decreased significantly during most of the rewarming period.



Figs. 3 and 4. Mean norepinephrine and epinephrine, respectively, at baseline (B) and during the 160-min rewarming period.






Fig. 5. Mean heart rate in beats per minute at baseline (B) and during the 160-min rewarming period.

This decrease in heart rate normally observed with immersion hypothermia was reduced by the administration of 3.5 mg/kg of caffeine immediately after subjects were removed from the cold water immersion bath.

#### DISCUSSION

These data demonstrate that subjects made mildly hypothermic after immersion in cold water exhibit a smaller afterdrop when given oral caffeine immediately upon removal from cold water. The lack of any differences in the skin temperature and plasma norepinephrine levels suggests that the effect of caffeine on

rewarming was not because of altered blood flow in the superficial skin. However, it is possible that skin thermistors utilized in the present study may not have been sensitive to subtle changes in vasomotor tone or sensitive to blood flow changes at deeper skin levels where caffeine could possibly have modified vascular flow (12).

As the rewarming slopes between the caffeine and control conditions were similar during the initial and latter part of the rewarming period, we believe that caffeine truncated the magnitude of the afterdrop without necessarily altering the rate of recovery. Since a direct influence on blood vessel vaso-reactivity appears unlikely to account for the observed differences, it is probable that the effect of caffeine on the afterdrop resulted from a stimulatory effect on metabolic rate or from caffeine maintaining a stable metabolic rate under conditions in which it would normally decrease (13). In this regard, studies have shown that the methylxanthines increase basal metabolism and can indeed improve cold tolerance (14). Although metabolic rate was not explicitly measured in this experiment, the sustained high levels of plasma epinephrine levels, together with a similar pattern observed in the heart rate after caffeine administration, suggest that caffeine may have induced a slight enhancement of metabolic rate during the rewarming period (13) above the level normally seen during rewarming from mild hypothermia. Such a metabolic increase, especially in the initial period following removal from cold



immersion, could have reduced the amount of afterdrop. That caffeine administration increased systemic epinephrine levels and heart rate is consistent with its mechanism of action that has been reported previously (4, 5, 6, 9, 10). The lack of a significant effect with plasma norepinephrine indicates that while caffeine is a potent stimulus to noradrenergic release (4, 6, 10), its effects are likely to be overshadowed by the profound effects of cold on peripheral circulation (15).

In summary, caffeine attenuated the magnitude of the post-immersion afterdrop normally seen upon a subject's removal from cold water. Although the mechanism for this effect is not completely understood, it is likely that events which stimulate oxygen consumption directly and influence internal heat balance are most prominent. That caffeine and possibly other methylxanthine compounds, given orally, can facilitate rewarming after hypothermia suggests that their administration, together with other rewarming methods (16, 17, 18, 19), might prove to be a potentially useful adjunctive treatment to increase patient survival from cold water immersion hypothermia.

## REFERENCES

1. Lonning, P., A. Skulberg, and F. Abyholm. Accidental hypothermia. *Acta Anesthesiol. Scand.* 1986; 301:601-613.
2. Webb, P. Afterdrop of body temperature during rewarming: an alternative explanation. *J. Appl. Physiol.* 1986; 60(2):385-390.
3. Romet, T. T. Mechanism of afterdrop after cold water immersion. *J. Appl. Physiol.* 1988; 65(4):1535-1538.
4. Lane, J. D. and R. B. Williams. Cardiovascular effects of caffeine and stress in regular coffee drinkers. *Psychophysiology.* 1987; 4(2):157-164.
5. Pincomb, G. A., W. R. Lovallo, R. B. Passey, T. L. Whitsett, S. M. Silverstein, and M. F. Wilson. Effects of caffeine on vascular resistance, cardiac output and myocardial contractility in young men. *Amer. J. Cardiol.* 1985; 56:119-122.
6. Robertson, D., J. C. Frolich, R. K. Carr, J. T. Watson, J. W. Hollifield, D. G. Shand, and J. A. Oates. Effects of caffeine on plasma renin activity, catecholamines and blood pressure. *New Engl. J. Med.* 1978; 298:181-186.
7. Zahn, T. P. and J. L. Rapoport. Autonomic nervous system effects of acute doses of caffeine in caffeine users and abstainers. *Int. J. Psychophysiology.* 1987; 5:33-41.
8. Von Borstel, W. R. and R. J. Wurtman. Caffeine and the cardiovascular effects of physiological levels of adenosine.

- In: *Caffeine: Perspectives from Recent Research*. P. B. Dews (Ed), Springer - Verlag, New York, 1984; 142-150.
9. Bellett, S., L. Roman, O. DeCastro, K. E. Kim, and A. Kershbaum. Effect of coffee ingestion on catecholamine release. *Metabolism*. 1969; 18:288-291.
  10. Fernstrom, J. D. and M. H. Fernstrom. Effects of caffeine on monoamine neurotransmitters in the central and peripheral nervous system. In: *Caffeine: Perspectives from Recent Research*. P. B. Dews (Ed), Springer - Verlag, New York, 1984; 107-118.
  11. DuBois, D. and E. F. DuBois. Clinical calorimetry. A formula to estimate the approximate surface area if height and weight be known. *Arch. Inter. Med.* 1916; 17:863-871.
  12. Nielsen, B. Thermoregulation in rest and exercise. *Acta Physiol. Scan. Suppl.* 1969; 323:1-74.
  13. Staten, M. A., D. E. Matthews, P. E. Cryer, and D. M. Bier. Physiological increments in epinephrine stimulate metabolic rate in humans. *Am. J. Physiol.* 1987; 253 (Endocrinol. Metab. 16):E322-E330.
  14. Wang, L. C. H. and E. C. Anholt. Elicitation of supramaximal thermogenesis by aminophylline in the rat. *J. Appl. Physiol.* 1982; 53(1):16-20.
  15. Savard, G. K., K. E. Cooper, W. L. Veale., and T. J. Malkinson. Peripheral blood flow during rewarming from mild hypothermia in humans. *J. Appl. Physiol.* 1985; 58(1):4-13.

16. Giesbrecht, G. G., G. K. Bristow, A. Uin, A. E. Ready, and R. A. Jones. Effectiveness of three field treatments for induced mild (33.0°C) hypothermia. *J. Appl. Physiol.* 1987; 63(6):2375-2379.
17. Golden, F. St. C. Rewarming. In: *The Nature and Treatment of Hypothermia*. R. S. Pozos and L. E. Wittmers (Eds), University of Minnesota Press, Minneapolis, 1983; 194-208.
18. Harnett, R. M., J. R. Pruitt, and F. R. Sias. A review of the literature concerning resuscitation from hypothermia: part II - selected rewarming protocols. *Aviat. Space Environ. Med.* 1983; 54(6):487-495.
19. Hesslink, R., S. Pepper, R. Olsen, S. B. Lewis, and L. D. Homer. Radio frequency (13.56 MHz) energy enhances recovery from mild hypothermia. *J. Appl. Physiol.* 1989; 67(3):1208-1212.